

ANTIFEN 2 HC

Food for Special Medical Purposes

Suitable for dietary management of subjects affected by phenylketonuria and hyperphenylalaninemia
FROM THREE YEARS



Net weight: 500 g

Food for special medical purposes with sugars and sweeteners

INGREDIENTS: L-leucine, L-tyrosine, L-aspartic acid, Glycine, L-lysine hydrochloride, Calcium glycerol phosphate, L-Glutamine, L-Proline, L-Valine, flavors, L-arginine, L-threonine, L-isoleucine, L-serine, Potassium citrate, L-alanine, L-histidine, Magnesium citrate, Thickener: xanthan gum E415; Sodium citrate, L-methionine, L-tryptophan, Choline bitartrate, Sweeteners: acesulfame k E950, sucralose E955; Sucrose, Sodium chloride, L-cystine, Calcium citrate, Dried glucose syrup, Corn maltodextrin, Taurine, Ferrous bisglycinate, Vitamin C (L-ascorbic acid), L-carnitine L-tartrate, Vitamin E (DL alpha tocopherol acetate), Zinc chloride, Niacin (nicotinamide), Manganese gluconate, Cupric gluconate, Pantothenic acid (D-pantothenate calcium), Vitamin B6 (pyridoxine hydrochloride), Vitamin B2 (riboflavin), Vitamin B1 (thiamin hydrochloride), Sodium molybdate, Vitamin A (retinyl palmitate), Folic acid (pteroylmonoglutamic acid), Chromium picolinate, Potassium iodide, Sodium selenite, Vitamin K1 (phyloquinone), D-biotin, Vitamin D3 (cholecalciferol), Vitamin B12 (cyanocobalamin).

INDICATION: Suitable for dietary management of subjects affected by phenylketonuria and hyperphenylalaninemia from three years old. Powdered phenylalanine free protein substitute, containing essential and non-essential amino acids, carbohydrate, vitamins, minerals and trace elements. With sugar and sweeteners.

IMPORTANT NOTICE: The product must be prescribed and used under medical control and it is intended for subjects aged three or above. It is not suitable for use as the sole source of nourishment and it is not suitable for parenteral use. The product should not be given to healthy children and adults.

DOSE AND INSTRUCTION FOR USE: The daily dose of the product must be determined by the physician and is dependent on the age, body weight and individual tolerance to phenylalanine. As standard dilution dissolve: 33 g in 200 ml of water: 20 g PE. 1 measuring cup = 8,3 g of powder. pour the water into a large and well-cleaned container, add the prescribed dose of powder and shake well until a homogeneous dispersion is obtained. The reconstituted product can be stored for a few hours. Throw away the unused product. Shake well before use.

STORAGE: Store in a cool dry place away from light and moisture. The expiry date refers to the unopened product, correctly stored. Once opened, use within 30 days.

Nutrients, per	100 g				
Energy	kJ	1129	L-Valine	g	5
Energy	kcal	270	L-Carnitine	mg	60
Fat	g	0,5	L-Taurine	mg	160
of which saturates	g	0,2	Vitamins		
monounsaturates	g	0	Vitamin A	µg	1100
polyunsaturates	g	0	Vitamin D	µg	30
Linoleic acid	mg	0	Vitamin E	mg	20
α-linolenic acid	mg	0	Vitamin K	µg	110
DHA	mg	0	Thiamin (B1)	mg	1,9
ARA	mg	0	Riboflavin (B2)	mg	2,1
Carbohydrates	g	4,9	Niacin (B3)	mg	24
of which sugars	g	0,7	Panthenic ac.	mg	7,8
GOS	g	0	Vitamin B6	mg	2,2
FOS	g	0	Biotin	µg	60
Protein Equivalent	g	60	Folate	µg	540
Salt	g	2	Vitamin B12	µg	5,6
Aminoacids			Vitamin C	mg	110
L-Alanine	g	2,6	Choline	mg	410
L-Arginine	g	4,4	Inositol	mg	0
L-Aspartic ac.	g	6,6	Minerals		
L-Cystine	g	0,6	Sodium	mg	800
L-Phenylalanine	g		Calcium	mg	1200
L-Glutamine	g	5,3	Potassium	mg	1100
Glycine	g	6,6	Phosphorus	mg	850
L-Histidine	g	2,6	Chloride	mg	500
L-Isoleucine	g	3,8	Magnesium	mg	350
L-Leucine	g	7,5	Trace elements		
L-Lysine	g	5,3	Iron	mg	22
L-Methionine	g	1,5	Zinc	mg	18
L-Proline	g	5,1	Manganese	mg	3,1
L-Serine	g	3,2	Copper	mg	1,6
L-Threonine	g	4,2	Iodine	µg	210
L-Tyrosine	g	6,8	Molybdenum	µg	60
L-Triphophan	g	1,4	Chromium	µg	40
			Selenium	µg	90