

ANTIFEN 3 HC

Food for Special Medical Purposes

Suitable for dietary management of subjects affected by phenylketonuria and hyperphenylalaninemia
FROM TEN YEARS



Net weight: 20x 28,6 g – 572 g

Food for special medical purposes with sweeteners

INGREDIENTS: L-Glutamine, L-Leucine, L-Tyrosine, L-Lysine hydrochloride, L-Aspartic Acid, Glycine, L-Proline, L-Valine, Calcium glycerol phosphate, L-Arginine, L-Threonine, L-Isoleucine, L-Serine, Flavor, L-Alanine, L-Histidine, Potassium Citrate, Magnesium Citrate, L-Methionine, L-Tryptophan, Sodium Citrate, Corn maltodextrin, Choline Bitartrate, Sodium Chloride, L-Cystine, Calcium Citrate, Inositol, Taurine, Ferrous Bysglicinate, Vitamin C (L-Ascorbic Acid), L-Carnitine L-Tartrate, Sweeteners: Acesulfame K E950, Sucralose E955; Vitamin E (DL-alpha tocopherol acetate), Zinc Chloride, Niacin (Nicotinamide), Manganese Gluconate, Vitamin D (Cholecalciferol), Cupric Gluconate, Pantothenic acid (D-Pantothenate calcium), Vitamin A (Retinyl Acetate), Vitamin B6 (Pyridoxine Hydrochloride), Vitamin K (Phylloquinone), Vitamin B1 (Thiamin Hydrochloride), Vitamin B2 (Riboflavin), Sodium Molybdate, Folic Acid (Pteroylmonoglutamic acid), Chromium Picolinate, Potassium Iodide, Sodium Selenite, D-Biotin, Vitamin B12 (Cyanocobalamin).

INDICATION: Suitable for dietary management of subjects affected by phenylketonuria and hyperphenylalaninemia from ten years old. Powdered phenylalanine free protein substitute, containing essential and non-essential amino acids, vitamins, minerals and trace elements. With sweeteners.

IMPORTANT NOTICE: The product must be prescribed and used under medical control and it is intended for subjects aged ten or above. It is not suitable for use as the sole source of nutrition and it is not suitable for parenteral use. The product should not be given to healthy children and adults. Keep out of reach of children. Do not take/administer in case of allergy and/or known or presumed intolerance to one or more of the ingredients.

DOSE AND INSTRUCTION FOR USE: The daily dose of the product must be determined by the physician and is dependent on the age, body weight and individual tolerance to phenylalanine. As standard dilution dissolve: 16.6 g in 100 ml of water: 12 g PE. 1 measuring cup = 8,3 g of powder. Pour the water into a large and well-cleaned container, add the prescribed dose of powder and shake well until a homogeneous dispersion is obtained. The reconstituted product can be stored for a few hours. Throw away the unused product. Shake well before use.

STORAGE: Store in a cool dry place away from light and moisture. The expiry date refers to the unopened product, correctly stored. Once opened, use within 30 days.

Nutrients, per		100 g		
Energy	kJ	1206/288	L-Valine	g 5,4
Energy	kcal	0	L-Carnitine	mg 65
Fat	g	0	L-Taurine	mg 174
of which saturates	g	0	Vitamins	
monounsaturates	g	0	Vitamin A	µg 990
polyunsaturates	g	0	Vitamin D	µg 27
Linoleic acid	mg	0	Vitamin E	mg 18
α-linolenic acid	mg	0	Vitamin K	µg 99
DHA	mg	0	Thiamin (B1)	mg 1,7
ARA	mg	2	Riboflavin (B2)	mg 1,9
Carbohydrates	g	1	Niacin (B3)	mg 21,6
of which sugars	g	0	Panthenic ac.	mg 7
GOS	g	0	Vitamin B6	mg 2
FOS	g	0	Biotin	µg 54
Protein Equivalent	g	70	Folate	µg 486
Salt	g	1,5	Vitamin B12	µg 5
Aminoacids			Vitamin C	mg 99
L-Alanine	g	2,8	Choline	mg 369
L-Arginine	g	4,8	Inositol	mg 260
L-Aspartic ac.	g	7,2	Minerals	
L-Cystine	g	0,7	Sodium	mg 611
L-Phenylalanine	g		Calcium	mg 1080
L-Glutamine	g	10,9	Potassium	mg 990
Glycine	g	7,2	Phosphorus	mg 765
L-Histidine	g	2,8	Chloride	mg 450
L-Isoleucine	g	4,1	Magnesium	mg 315
L-Leucine	g	8,2	Trace elements	
L-Lysine	g	5,8	Iron	mg 19,8
L-Methionine	g	1,6	Zinc	mg 16,2
L-Proline	g	5,6	Manganese	mg 2,8
L-Serine	g	3,5	Copper	mg 1,4
L-Threonine	g	4,6	Iodine	µg 189
L-Tyrosine	g	7,4	Molybdenum	µg 54
L-Triphophan	g	1,5	Chromium	µg 36
			Selenium	µg 81