

ANTIFEN 3 LC VANILLA

Food for Special Medical Purposes

Suitable for dietary management of subjects affected by phenylketonuria and hyperphenylalaninemia
FROM TEN YEARS



Net weight: 20x 50 g – 1 kg

Food for special medical purposes with sugars and sweeteners

INGREDIENTS: Corn maltodextrin, Dried glucose syrup, Sucrose, L-Glutamine, L-Leucine, Calcium glycerol phosphate, L-Tyrosine, L-Aspartic Acid, Glycine, L-Lysine hydrochloride, Flavor, L-Proline, L-Valine, L-Arginine, L-Threonine, L-Isoleucine, Potassium Citrate, L-Serine, Magnesium Citrate, Inulin, L-Alanine, L-Histidine, Sodium Citrate, L-Methionine, L-Tryptophan, Choline Bitartrate, Sodium Chloride, L-Cystine, Calcium Citrate, Inositol, Ferrous Bysglycinate, Vitamin C (L-Ascorbic Acid), Taurine, Sweeteners: Acesulfame K E950, Sucralose E955; L-Carnitine L-Tartrate, Vitamin E (DL-alpha tocopherol acetate), Zinc Chloride, Niacin (Nicotinamide), Manganese Gluconate, Vitamin D (Cholecalciferol), Cupric Gluconate, Pantothenic acid (D-Pantothenate calcium), Vitamin A (Retenyl Acetate), Vitamin B6 (Pyridoxine Hydrochloride), Vitamin B2 (Riboflavin), Vitamin B1 (Thiamin Hydrochloride), Vitamin K (Phylloquinone), Sodium Molybdate, Folic Acid (Pteroylmonoglutamic acid), Chromium Picolinate, Potassium Iodide, Sodium Selenite, D-Biotin, Vitamin B12 (Cyanocobalamin).

INDICATION: Suitable for dietary management of subjects affected by phenylketonuria and hyperphenylalaninemia from ten years old. Vanilla Flavor. Powdered phenylalanine free protein substitute, containing essential and non-essential amino acids, carbohydrate, vitamins, minerals and trace elements. With sugars and sweeteners.

IMPORTANT NOTICE: The product must be prescribed and used under medical control and it is intended for subjects aged ten or above. It is not suitable for use as the sole source of nutrition and it is not suitable for parenteral use. The product should not be given to healthy children and adults. Keep out of reach of children. Do not take/administer in case of allergy and/or known or presumed intolerance to one or more of the ingredients.

DOSE AND INSTRUCTION FOR USE: The daily dose of the product must be determined by the physician and is dependent on the age, body weight and individual tolerance to phenylalanine. As standard dilution dissolve: 50 g in 200 ml of water: 20 g PE. 1 measuring cup = 8,3 g of powder. pour the water into a large and well-cleaned container, add the prescribed dose of powder and shake well until a homogeneous dispersion is obtained. The reconstituted product can be stored for a few hours. Throw away the unused product. Shake well before use.

STORAGE: Store in a cool dry place away from light and moisture. The expiry date refers to the unopened product, correctly stored. Once opened, use within 30 days.

Nutrients, per		100 g		
Energy	kJ	1293/309	L-Valine	g 3,1
Energy	kcal	0	L-Carnitine	mg 38
Fat	g	0	L-Taurine	mg 100
of which saturates	g	0	Vitamins	
monounsaturates	g	0	Vitamin A	µg 825
polyunsaturates	g	0	Vitamin D	µg 22,5
Linoleic acid	mg	0	Vitamin E	mg 15
α-linolenic acid	mg	0	Vitamin K	µg 62,5
DHA	mg	0	Thiamin (B1)	mg 1,4
ARA	mg	36,5	Riboflavin (B2)	mg 1,6
Carbohydrates	g	9,6	Niacin (B3)	mg 18
of which sugars	g	0	Pantothenic ac.	mg 5,9
GOS	g	0	Vitamin B6	mg 1,7
FOS	g	1,5	Biotin	µg 45
Protein Equivalent	g	40	Folate	µg 405
Salt	g	1,5	Vitamin B12	µg 4,2
Aminoacids			Vitamin C	mg 82,5
L-Alanine	g	1,6	Choline	mg 308
L-Arginine	g	2,8	Inositol	mg 150
L-Aspartic ac.	g	4,1	Minerals	
L-Cystine	g	0,4	Sodium	mg 600
L-Phenylalanine	g	0	Calcium	mg 900
L-Glutamine	g	6	Potassium	mg 825
Glycine	g	4,1	Phosphorus	mg 637
L-Histidine	g	1,6	Chloride	mg 375
L-Isoleucine	g	2,4	Magnesium	mg 262
L-Leucine	g	4,7	Trace elements	
L-Lysine	g	3,3	Iron	mg 16,5
L-Methionine	g	0,9	Zinc	mg 13,5
L-Proline	g	3,2	Manganese	mg 2,3
L-Serine	g	2	Copper	mg 1,2
L-Threonine	g	2,6	Iodine	µg 157
L-Tyrosine	g	4,3	Molybdenum	µg 45
L-Triphophan	g	0,9	Chromium	µg 30
			Selenium	µg 68