


GLYTACTIN[™] BUILD RASPBERRY LEMONADE FLAVOUR

Food for special medical purposes (FSMP)
For the dietary management of Phenylketonuria (PKU) and hyperphenylalaninemia (HPA)
FROM 3 YEARS



NET WT: 621 g

Food for special medical purposes (FSMP). With sweetener.

Protein Powder with Vitamins and Minerals

5 mg Phe per 5 g Protein Equivalent

INGREDIENTS: Glycomacropeptide (milk), L-arginine, L-leucine, L-tyrosine, anti-foaming agent (soy lecithin), glycine, L-lysine acetate, sunflower oil, contains less than 2% of L-histidine, L-tryptophan, DHA algal oil (derived from microalgae Schizochytrium sp.), maltodextrin, food starch modified, inulin, thickeners (guar gum, xanthan gum, carrageenan), salt, natural flavours, vitamin A palmitate, thiamin mononitrate, riboflavin, pyridoxine hydrochloride, D-calcium pantothenate, niacinamide, phytonadione, menaquinone-MK7, folic acid, biotin, cyanocobalamin, ascorbic acid, cholecalciferol, DL- α -tocopheryl acetate, calcium lactate, potassium phosphate, magnesium phosphate, ferrous fumarate, zinc sulphate, manganese sulphate, copper gluconate, potassium iodide, sodium selenite, sodium molybdate, chromium chloride, potassium chloride, canola oil, sweetener (sucralose), acidulant (citric acid), colour (beetroot red), Bacillus coagulans GBI-30 6086. Contains milk and soy.

INDICATIONS: For the Dietary Management of Phenylketonuria/Hyperphenylalaninemia. GLYTACTIN BUILD is a low-phenylalanine mixture based on glycomacropeptide (GMP), amino acids, vitamins, minerals, trace elements with docosahexaenoic acid (DHA).

IMPORTANT NOTICE: Must be administered under medical supervision only. Not suitable for use as a sole source of nourishment. Protein in prescribed amounts must be supplemented to completely meet phenylalanine requirements. For enteral use only. Not for parenteral use. Suitable for individuals with PKU over 3 years of age.

DIRECTIONS FOR USE: Use as directed by a physician. Must be administered under medical supervision only. Recommended dilution: 1 heaping scoop (approximately 8g powder) mixed with 30mL of water. For best results, weigh powder on a gram scale. Dilution may be adjusted to taste. Mixing recommendation: measure water in container, add prescribed amount of powder, and shake well until fully blended. Once prepared, product can be refrigerated for up to 24 hours. Shake well prior to use. Store pouch in a cool, dry place. Avoid extreme temperatures. Once pouch is opened, reseal.

Nutrients, per			8 g scoop			100 g		
	8 g scoop	100 g		mg	0,13		mg	1,6
Energy	kcal	28	350	Folate	μ g DFE	31	383	
Energy	kJ	117	1464	Folic Acid	μ g	18	225	
Fat	g	0,5	6,3	Vitamin B12	μ g	0,23	2,9	
Saturated Fat	g	0,1	0,7	Pantothenic Acid (B5)	mg	0,56	7,0	
Trans Fat	g	0,0	0,0	Biotin	μ g	2,5	31	
DHA	mg	13	164	Choline	mg	7,4	93	
Cholesterol	mg	0,1	0,8	Calcium	mg	139	1738	
Carbohydrates	g	0,4	5,3	Chromium	μ g	4,0	50	
Sugar	g	0,1	0,8	Copper	μ g	51	639	
Sugar Alcohols	g	0,0	0,0	Iodine	μ g	18	219	
Fibre	g	0,1	1,0	Iron	mg	2,0	25	
Protein Equivalent	g	5,2	64	Magnesium	mg	25	313	
Phenylalanine	mg	5,0	62	Manganese	mg	0,26	3,2	
Salt	g	0,17	2,1	Molybdenum	μ g	4,6	58	
Vitamin A	μ g RAE	84	1050	Phosphorus	mg	128	1604	
Vitamin C	mg	14	169	Selenium	μ g	6,1	77	
Vitamin D	μ g	3,7	46	Zinc	mg	1,1	14	
Vitamin E	mg α -TE	1,5	19	Potassium	mg	143	1783	
Vitamin K1	μ g	4,6	57	Sodium	mg	69	857	
Vitamin K2 (MK-7)	μ g	4,6	57	Chloride	mg	58	729	
Thiamin (B1)	mg	0,13	1,6	Inositol	mg	0,0	0,0	
Riboflavin (B2)	mg	0,13	1,6	Osmolality	mOsm/kg		898	
Niacin (B3)	mg NE	2,7	34	pH			6,0	